

## **I. AMENDMENTS TO THE CLAIMS**

This listing of claims will replace all prior versions, and listings, of claims in the application:

1-37. (Canceled)

38. (Currently Amended) A method for obtaining the production of a fertilizable oocyte within a program of controlled ovarian stimulation for assisted reproduction techniques (COS/ART) consisting essentially of:

- (a) administering a combination of luteinizing hormone (LH) and follicle stimulating hormone (FSH) to induce follicle growth, wherein the dose of LH and FSH remains the same throughout the treatment period, and
- (b) administering an amount of a luteinizing hormone releasing hormone (LHRH) antagonist selected from the group consisting of Ganirelix, Antarelix, Antide, Azaline B, Ramorelix, A-76154, Nal-Glu, 88-88, Cetrorelix, D23980, and D-24824, to prevent a premature LH surge, wherein the LHRH antagonist is administered in a single or dual dosage regimen of 3 mg per dose beginning on menstruation cycle day 1 to 10;

wherein follicular growth occurs in the absence of a LH surge, a fertilizable oocyte is produced, ovulation occurs between day 9 and 20 of the menstruation cycle,

~~and the dosage regimen of the LHRH antagonist is selected so as to suppress~~ wherein said amount of LHRH antagonist suppresses endogenous LH secretion,

~~[[but]] wherein said amount of LHRH antagonist~~ does not suppress endogenous FSH secretion, which is maintained at a natural level and individual estrogen development is not affected.

39. (Previously presented) The method of claim 38, wherein step (a) comprises administering human menopausal gonadotropin (HMG) to induce follicle growth.

40-41. (Canceled)

42. (Previously Presented) The method of claim 38, wherein the LHRH antagonist is administered by subcutaneous injection.

43. (Canceled)

44. (Previously presented) The method of claim 38, wherein the LHRH antagonist is administered starting on cycle day 4 to 8.

45. (Previously Presented) The method of claim 38, wherein the LHRH antagonist is administered starting on cycle day 6 to 10 and ovulation occurs between day 9-16 of the menstruation cycle.

46. (Previously Presented) The method of claim 38, wherein ovulation occurs within 6.5 days following administration of a single or second dose of the LHRH antagonist.

47. (Previously Presented) The method of claim 38, wherein ovulation occurs normally, without the administration of a hormone or hormone agonist to induce ovulation.

48. (Previously Presented) The method of claim 38, wherein ovulation is induced by administering a hormone or hormone agonist in order to induce ovulation.

49. (Previously Presented) The method of claim 38, wherein ovulation is induced by administering a hormone or hormone agonist selected from the group consisting of native LH, recombinant LH, an LHRH agonist, and HCG.

50. (Previously Presented) The method of claim 38, wherein the LHRH antagonist is Cetrorelix.

51. (Currently Amended) A method for obtaining the production of a fertilizable oocyte within a program of COS/ART consisting essentially of:

(a) administering human menopausal gonadotropin (HMG) to induce follicle growth, wherein the dose of HMG remains the same throughout the treatment period, and

(b) administering an amount of Cetorelix to prevent a premature LH surge, wherein Cetorelix is administered in a single or dual dosage regimen of 3 mg per dose beginning on menstruation cycle day 1 to 10;

wherein follicular growth occurs in the absence of a LH surge, a fertilizable oocyte is produced, ovulation occurs between day 9 and 20 of the menstruation cycle,

~~and the dosage regimen of the wherein said amount of Cetorelix is selected so as to suppress suppresses~~ endogenous LH secretion,

~~[[but]]~~ wherein said amount of Cetorelix does not suppress endogenous FSH secretion, which is maintained at a natural level and individual estrogen development is not affected.

52-55. (Canceled)

56. (Previously Presented) The method of claim 51, wherein the Cetorelix is administered starting on cycle day 4 to 8.

57. (Previously Presented) The method of claim 51, wherein Cetorelix is administered starting on cycle day 6 to 10 and ovulation occurs between day 9-16 of the menstruation cycle.

58. (Previously Presented) The method of claim 51, wherein ovulation occurs within 6.5 days following administration of a single or second dose of Cetorelix.

59. (Previously Presented) The method of claim 51, wherein ovulation occurs normally, without the administration of a hormone or hormone agonist to induce ovulation.

60. (Previously Presented) The method of claim 51, wherein ovulation is induced by administering a hormone or hormone agonist selected from the group consisting of native LH, recombinant LH, an LHRH agonist, and HCG.

61. (Currently Amended) A method for obtaining the production of a fertilizable oocyte within a program of COS/ART comprising:

(a) administering a combination of luteinizing hormone (LH) and follicle stimulating hormone (FSH) to induce follicle growth; and

(b) administering an amount of an LHRH antagonist selected from the group consisting of Ganirelix, Antarelix, Antide, Azaline B, Ramorelix, A-76154, Nal-Glu, 88-88, Cetorelix, D-23980, and D-24824 to prevent a premature LH surge;

wherein the LHRH antagonist is administered in a single or dual dosage regimen of 1 to 10 mg per dose beginning on menstruation cycle day 1 to 10;

wherein follicular growth occurs in the absence of a LH surge, a fertilizable oocyte is produced, ovulation occurs normally between day 9 and 20 of the menstruation cycle without the administration of a hormone or hormone agonist to induce ovulation,

~~and the~~ wherein said amount of LHRH antagonist is sufficient to suppress LH,

~~while~~ wherein said amount of LHRH antagonist does not suppress endogenous FSH secretion which is maintained at a natural level and individual estrogen development is not affected.

62. (Previously Presented) The method of claim 61, wherein the dosage of LHRH antagonist is in the range of 2-6 mg per dose.

63. (Previously Presented) The method of claim 61, wherein the dosage of LHRH antagonist is 3 mg per dose.

64. (Canceled)

65. (Previously Presented) The method of claim 61, wherein the LHRH antagonist is administered by subcutaneous injection.

66. (Canceled)

67. (Previously Presented) The method of claim 61, wherein the LHRH antagonist is administered starting on cycle day 4 to 8.

68. (Previously Presented) The method of claim 61, wherein the LHRH antagonist is administered starting on cycle day 6 to 10 and ovulation occurs between day 9-16 of the menstruation cycle.

69. (Previously Presented) The method of claim 61, wherein ovulation occurs within 6.5 days following administration of a single or second dose of the LHRH antagonist.

70. (Previously Presented) The method of claim 61, wherein step (a) comprises administering human menopausal gonadotropin (HMG) to induce follicle growth.

71. (Canceled)

72. (Previously Presented) The method of claim 61, wherein the LHRH antagonist is Cetrorelix.

73. (Previously Presented) The method of claim 70, wherein the LHRH antagonist is Cetrorelix.

74. (Previously Presented) The method of claim 73, wherein the dosage of Cetrorelix is in the range of 2-6 mg per dose.

75. (Previously Presented) The method of claim 73, wherein the dosage of LHRH antagonist is 3 mg per dose.

76-77. (Canceled)

78. (Previously Presented) The method of claim 73, wherein the LHRH antagonist is administered starting on cycle day 4 to 8.

79. (Previously Presented) The method of claim 73, wherein Cetrorelix is administered starting on cycle day 6 to 10 and ovulation occurs between day 9-16 of the menstruation cycle.

80. (Previously Presented) The method of claim 73, wherein ovulation occurs within 6.5 days following administration of a single or second dose of Cetrorelix.

81-82. (Canceled)

83. (Currently amended) A method for obtaining the production of a fertilizable oocyte within a program of controlled ovarian stimulation for assisted reproduction techniques (COS/ART) comprising

(a) administering a combination of luteinizing hormone (LH) and follicle stimulating hormone (FSH) to induce follicle growth,

(b) administering an amount of an LHRH antagonist selected from the group consisting of Ganirelix, Antarelix, Antide, Azaline B, Ramorelix, A-76154, Nal-Glu, 88-88, Cetrorelix, D-23980, and D-24824 to prevent a premature LH surge, wherein the LHRH antagonist is administered in a dosage regimen of daily doses of 0.25 mg/day for multiple days,

wherein the LHRH antagonist is administered daily beginning on menstruation cycle day 1 to 10, wherein the follicular growth occurs in the absence of a LH surge, a fertilizable oocyte is produced, ovulation occurs between day 9 and 20 of the menstruation cycle,

~~and the wherein said amount of~~ LHRH antagonist is sufficient to suppress LH,

~~while wherein said amount of LHRH antagonist does not suppress endogenous~~ FSH secretion ~~which~~ is maintained at a natural level and individual estrogen development is not affected.

84. (Previously Presented) The method of claim 83, wherein the LHRH antagonist is administered by subcutaneous injection.

85. (Canceled)

86. (Previously Presented) The method of claim 83, wherein the LHRH antagonist is administered starting cycle on day 4 to 8.

87. (Previously Presented) The method of claim 83, wherein a daily dose of the LHRH antagonist is administered for 3 to 14 days.

88. (Previously Presented) The method of claim 83, wherein a daily dose of the LHRH antagonist is administered for 3 to 7 days.

89. (Previously Presented) The method of claim 83, wherein ovulation occurs normally, without the administration of a hormone or hormone agonist to induce ovulation.

90. (Previously Presented) The method of claim 83, wherein ovulation is induced by administering a hormone or hormone agonist selected from the group consisting of native LH, recombinant LH, an LHRH agonist, and HCG.

91. (Previously Presented) The method of claim 83, wherein the LHRH antagonist is Cetrorelix.

92. (Currently Amended) A method for obtaining the production of a fertilizable oocyte within a program of COS/ART comprising:

(a) administering human menopausal gonadotropin (HMG) to induce follicle growth, and;

(b) administering an amount of Cetrorelix to prevent a premature LH surge, wherein Cetrorelix is subcutaneously administered in a dosage regimen of daily doses of 0.25 mg per day for multiple days;

wherein follicular growth occurs in the absence of a LH surge, a fertilizable oocyte is produced, ovulation occurs between day 9 and 20 of the menstruation cycle,

~~and the~~ wherein said amount of Cetrorelix is sufficient to suppress LH,

~~while~~ and wherein FSH secretion is maintained at a natural level and individual estrogen development is not affected.

93. (Canceled)

94. (Previously Presented) The method of claim 92, wherein Cetrorelix is administered starting on cycle day 4 to 8.

95. (Previously Presented) The method of claim 92, wherein a daily dose of Cetrorelix is administered for 3 to 14 days.

96. (Previously Presented) The method of claim 92, wherein a daily dose of Cetorelix is administered for 3 to 7 days.

97. (Previously Presented) The method of claim 92, wherein ovulation occurs normally, without the administration of a hormone or hormone agonist to induce ovulation.

98. (Previously Presented) The method of claim 92, wherein ovulation is induced by administering a hormone or hormone agonist selected from the group consisting of native LH, recombinant LH, an LHRH agonist, and HCG.

99. (Currently Amended) A method for obtaining the production of a fertilizable oocyte within a program of controlled ovarian stimulation for assisted reproduction techniques (COS/ART) comprising

(a) administering a combination of luteinizing hormone (LH) and follicle stimulating hormone (FSH) to induce follicle growth; and

(b) administering an amount of an LHRH antagonist selected from the group consisting of Ganirelix, Antarelix, Antide, Azaline B, Ramorelix, A-76154, Nal-Glu, 88-88, Cetorelix, D-23980, and D-24824 to prevent a premature LH surge, wherein the LHRH antagonist is administered in a dosage regimen of daily doses of from 0.25 to 0.5 mg per day for multiple days;

wherein follicular growth occurs in the absence of a LH surge, a fertilizable oocyte is produced, ovulation occurs normally between day 9 and 20 of the menstruation cycle without the administration of a hormone or hormone agonist to induce ovulation,

~~and the~~ wherein said amount of said LHRH antagonist is sufficient to suppress LH,

~~while~~ wherein said amount of said LHRH antagonist does not suppress endogenous FSH secretion, which is maintained at a natural level and individual estrogen development is not affected.

100. (Previously Presented) The method of claim 99, wherein the LHRH antagonist is administered by subcutaneous injection.



101. (Canceled)

102. (Previously Presented) The method of claim 99, wherein the LHRH antagonist is administered starting cycle day 4 to 8.

103. (Previously Presented) The method of claim 99, wherein a daily dose of the LHRH antagonist is administered for 3 to 14 days.

104. (Previously Presented) The method of claim 99, wherein a daily dose of the LHRH antagonist is administered for 3 to 7 days.

105. (Previously Presented) The method of claim 99, wherein step (a) comprises administering human menopausal gonadotropin (HMG) to induce follicle growth.

106. (Canceled)

107. (Previously Presented) The method of claim 99, wherein the LHRH antagonist is Cetorelix.

108. (Previously Presented) The method of claim 105, wherein the LHRH antagonist is Cetorelix.

109. (Canceled)

110. (Previously Presented) The method of claim 108, wherein Cetorelix is administered starting on cycle day 4 to 8.

111. (Previously Presented) The method of claim 108, wherein a daily dose of Cetorelix is administered for 3 to 14 days.

112. (Previously Presented) The method of claim 108, wherein a daily dose of Cetorelix is administered for 3 to 7 days.

113. (Previously Presented) The method of claim 108, wherein ovulation occurs normally, without the administration of a hormone or hormone agonist to induce ovulation.

114. (Previously Presented) The method of claim 108, wherein ovulation is induced by administering a hormone or hormone agonist selected from the group consisting of native LH, and recombinant LH.

115. (Currently amended) A method for obtaining the production of a fertilizable oocyte within a program of assisted reproduction techniques comprising:

(a) allowing normal follicular growth and development to proceed in the absence of stimulation by an exogenous gonadotropin;

(b) administering an amount of an LHRH antagonist selected from the group consisting of Ganirelix, Antarelix, Antide, Azaline B, Ramorelix, A-76154, Nal-Glu, 88-88, Cetrorelix, D-23980, and D-24824, in a dosage regimen that prevents a premature LH surge, beginning on menstruation cycle day 1 to 10;

wherein follicular growth and development proceeds in the absence of a LH surge and a fertilizable oocyte is produced, ovulation occurs between day 9 and 20 of the menstruation cycle,

~~and the wherein said amount of said~~ LHRH antagonist is sufficient to suppress LH,

~~while wherein said amount of said LHRH antagonist does not suppress endogenous FSH secretion, which~~ is maintained at a natural level and individual estrogen development is not affected.

116. (Previously Presented) The method of claim 115, wherein the LHRH antagonist is administered by subcutaneous injection.

117. (Canceled)

118. (Previously Presented) The method of claim 115, wherein the LHRH antagonist is administered starting on cycle day 4 to 8.

119. (Previously Presented) The method of claim 115, wherein the method is performed on a patient in whom follicular growth is inadequate due to previous treatment with an LHRH antagonist, and step (a) comprises allowing normal follicular growth and development

to proceed in the absence of treatment of the patient with an LHRH antagonist and in the absence of stimulation by an exogenous gonadotropin.

120. (Canceled)

121. (Previously Presented) The method of claim 115, wherein ovulation occurs normally, without the administration of a hormone or hormone agonist to induce ovulation.

122. (Previously Presented) The method of claim 115, wherein ovulation is induced by administering a hormone or hormone agonist in order to induce ovulation.

123. (Previously Presented) The method of claim 115, wherein ovulation is induced by administering a hormone or hormone agonist selected from the group consisting of native LH, recombinant LH, an LHRH agonist, and HCG.

124-125. (Canceled)

126. (Previously Presented) The method of claim 115, wherein the LHRH antagonist is Cetrorelix.

127. (Previously Presented) The method of claim 115, wherein a fertilizable oocyte is produced within a program of extracorporeal fertilization by sperm injection.

128. (Previously Presented) The method of claim 115, wherein a fertilizable oocyte is produced within a program of extracorporeal fertilization by *in vitro* fertilization.

129. (Previously Presented) The method of claim 115, wherein the LHRH antagonist is administered in a single or dual dosage regimen of 1 to 10 mg per dose.

130. (Previously Presented) The method of claim 129, wherein the dosage of LHRH antagonist is in the range of 2-6 mg per dose.

131. (Previously Presented) The method of claim 129, wherein the LHRH antagonist is administered starting on cycle day 6 to 10 and ovulation occurs between day 9 to 16 of the menstruation cycle.

132. (Previously Presented) The method of claim 129, wherein ovulation occurs within 6.5 days following administration of a single or second dose of the LHRH antagonist.

133. (Previously Presented) The method of claim 129, wherein the LHRH antagonist is Cetorelix.

134. (Previously Presented) The method of claim 115, wherein the LHRH antagonist is administered in a dosage regimen of daily doses of from 0.25 to 0.5 mg/day for multiple days.

135. (Previously Presented) The method of claim 134, wherein a daily dose of the LHRH antagonist is administered for 3 to 14 days.

136. (Previously Presented) The method of claim 134, wherein a daily dose of the LHRH antagonist is administered for 3 to 7 days.

137. (Previously Presented) The method of claim 135, wherein the LHRH antagonist is Cetorelix.

138. (Previously Presented) The method of claim 137, wherein the daily dose of Cetorelix is 0.25 mg/day or 0.5 mg/day.

139. (Previously Presented) The method of claim 83, wherein step (a) comprises administering human menopausal gonadotropin (HMG) to induce follicle growth.

140. (Previously Presented) The method of claim 99, wherein the LHRH antagonist is Cetorelix and the daily dose is 0.25 mg/day or 0.5 mg/day.

141. (Previously Presented) The method of claim 108, wherein the daily dose of Cetorelix is 0.25 mg/day or 0.5 mg/day.